

AD-A158 263

EXISTENCE OF EQUILIBRIUM PRICES FOR A SIMPLE PLANNING
MODEL(U) STANFORD UNIV CA SYSTEMS OPTIMIZATION LAB
H HU JUN 85 SOL-85-10 N00014-85-K-0343

1/1

UNCLASSIFIED

F/G 12/2


NL



END

FILED

DTIC



NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

AD-A158 263



Systems
Optimization
Laboratory

EXISTENCE OF EQUILIBRIUM PRICES FOR A
SIMPLE PLANNING MODEL

Hui Hu

TECHNICAL REPORT SOL 85-10

June 1985

DTIC FILE COPY

This document has been approved
for public release and sale; its
distribution is unlimited.

DTIC
ELECTE
AUG 20 1985
S D E

Department of Operations Research
Stanford University
Stanford, CA 94305

5 8 14 050

SYSTEMS OPTIMIZATION LABORATORY
DEPARTMENT OF OPERATIONS RESEARCH
STANFORD UNIVERSITY
STANFORD, CALIFORNIA 94305

EXISTENCE OF EQUILIBRIUM PRICES FOR A
SIMPLE PLANNING MODEL

Hui Hu

TECHNICAL REPORT SOL 85-10

June 1985

Research and reproduction of this report were partially supported by the Department of Energy Contract DE-AM03-76SF00326, PA# DE-AT03-76ER72018; National Science Foundation Grants DMS-8420623 and ECS-8312142; Office of Naval Research Contract N00014-85-K-0343.

Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do NOT necessarily reflect the views of the above sponsors.

Reproduction in whole or in part is permitted for any purposes of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

DTIC
ELECTE
S AUG 20 1985 D
E

EXISTENCE OF EQUILIBRIUM PRICES FOR A SIMPLE PLANNING MODEL

Hui Hu

Abstract

Consider parametric LP:

$$\begin{aligned} \min \quad & -\theta \\ \text{s.t.} \quad & AY + \theta(-d^0 + M\hat{\pi}) \geq b \\ & 0 \leq Y \leq K, \quad \theta \geq 0 \end{aligned} \tag{I}$$

where M is a positive definite (not necessarily symmetric) matrix, $K > 0$, $\hat{\pi}$ is a parameter, $\hat{\pi} \in S = \{\pi \geq 0: e\pi = 1\}$.

For each fixed $\hat{\pi} \in S$, we can solve (I) and it's dual problem and get optimal θ^* , Y^* , σ^* , π^* . The question is, is there a $\hat{\pi} \in S$ such that after solving the corresponding LP and normalizing the dual price π^* , it turns out that $\pi^*/e\pi^* = \hat{\pi}$? In this paper, we are going to show that under certain conditions, such a $\hat{\pi}$ does exist.

The economic interpretation of the above model will be given at the end of the paper.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



EXISTENCE OF EQUILIBRIUM PRICES FOR A SIMPLE PLANNING MODEL

Hui Hu

Notation:

$$\begin{aligned}
 & \min \quad (0, 0, \dots, 0, -1) \begin{bmatrix} Y \\ \theta \end{bmatrix} \\
 & \text{s.t.} \\
 (P\hat{\pi}): \quad & \begin{bmatrix} -I & 0 \\ A & -d^0 + M\hat{\pi} \end{bmatrix} \begin{bmatrix} Y \\ \theta \end{bmatrix} \geq \begin{bmatrix} -K \\ b \end{bmatrix} \\
 & Y \geq 0, \quad \theta \geq 0
 \end{aligned}$$

which is equivalent to:

$$\begin{aligned}
 & \min \quad -\theta \\
 & \text{s.t.} \\
 & AY + \theta(-d^0 + M\hat{\pi}) \geq b \\
 & 0 \leq Y \leq K, \theta \geq 0
 \end{aligned}$$

The corresponding dual problem is:

$$\begin{aligned}
 & \max \quad (\sigma, \pi) \begin{bmatrix} -K \\ b \end{bmatrix} \\
 & \text{s.t.} \\
 (D\hat{\pi}): \quad & (\sigma, \pi) \begin{bmatrix} -I & 0 \\ A & -d^0 + M\hat{\pi} \end{bmatrix} \leq (0, 0, \dots, 0, -1) \\
 & \sigma \geq 0, \pi \geq 0
 \end{aligned}$$

which is equivalent to:

$$\begin{aligned} \max \quad & -\sigma K + \pi b \\ \text{s.t.} \quad & \pi A \leq \sigma \\ & \pi(-d^0 + M\hat{\pi}) \leq -1 \\ & \pi \geq 0, \sigma \geq 0 \end{aligned}$$

Let $F(P\hat{\pi})$ and $F(D\hat{\pi})$ denote the feasible regions of $(P\hat{\pi})$ and $(D\hat{\pi})$ respectively. $S = \{\pi \geq 0 : e\pi = 1\}$.

Definition. Let $D \subseteq \mathbb{R}^n$, $U \subseteq \mathbb{R}^m$, a point to set map $f : D \rightarrow P(U)$ is upper hemicontinuous at $\bar{x} \in D$ if for all $x^k \rightarrow \bar{x}$ and $y^k \in f(x^k)$ such that $y^k \rightarrow \bar{y}$, we have $\bar{y} \in f(\bar{x})$. If f is upper hemicontinuous at all $x \in D$, f is called upper hemicontinuous.

Lemma 1. If $\{Y : AY \geq b, 0 \leq Y \leq K\} \neq \emptyset$ and for all $\hat{\pi} \in S$, $d^0 - M\hat{\pi} > 0$, then for any $\hat{\pi} \in S$, $(P\hat{\pi})$ and $(D\hat{\pi})$ have optimal solutions.

Proof. From the assumption we know there exists $\bar{Y} \in \{Y : AY \geq b, 0 \leq Y \leq K\}$, therefore

$$\begin{bmatrix} \bar{Y} \\ 0 \end{bmatrix} \in \left\{ \begin{bmatrix} Y \\ \theta \end{bmatrix} \geq 0 : \begin{bmatrix} -1 & 0 \\ A & -d^0 + M\hat{\pi} \end{bmatrix} \begin{bmatrix} Y \\ \theta \end{bmatrix} \geq \begin{bmatrix} -K \\ b \end{bmatrix} \right\}$$

which implies that for any $\hat{\pi} \in S$, $(P\hat{\pi})$ is feasible.

On the other hand, we know that $\{x \geq 0 : Ax \geq b\}$ is bounded if and only if $\{x \geq 0 : Ax \geq 0\} = \{0\}$.

$$\left\{ \begin{bmatrix} Y \\ \theta \end{bmatrix} \geq 0 : \begin{bmatrix} -I & 0 \\ A & -d^0 + M\hat{\pi} \end{bmatrix} \begin{bmatrix} Y \\ \theta \end{bmatrix} \geq \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\} \\ = \left\{ \begin{bmatrix} Y \\ \theta \end{bmatrix} > 0 : \begin{array}{l} Y = 0 \\ (-d^0 + M\hat{\pi}) \theta > 0 \end{array} \right\} = \{0\} ,$$

therefore for each $\hat{\pi} \in S$, $F(\hat{\pi})$ is bounded, therefore for all $\hat{\pi} \in S$, $(P\hat{\pi})$ and $(D\hat{\pi})$ have optimal solutions. Q.E.D.

Now define a point to set mapping f : For all $\hat{\pi} \in S$,

$$f(\hat{\pi}) = \{\pi : (\sigma, \pi) \text{ is optimal solution of } (D\hat{\pi})\} .$$

Under the assumption of Lemma 1, f is well defined on S , i.e., for all $\hat{\pi} \in S$, $f(\hat{\pi}) \neq \emptyset$, and furthermore, $f(\hat{\pi})$ is a convex set (that is because $f(\hat{\pi})$ is the projection of a convex set into a lower dimension).

Normalizing f , we get another point to set mapping \bar{f} : for all $\hat{\pi} \in S : \bar{f}(\hat{\pi}) = \{\pi / e\pi : \pi \in f(\hat{\pi})\}$.

It is easy to see that under the assumption of Lemma 1, for all $\hat{\pi} \in S$, $\bar{f}(\hat{\pi}) \neq \emptyset$, and $\bar{f}(\hat{\pi})$ is a convex subset of S .

Using these definitions, our question becomes: is there a fixed point of \bar{f} ?

Theorem 1. If for all $\hat{\pi} \in S$, $d^0 - M\hat{\pi} \geq \bar{d} > 0$ and $\{Y : AY > b, 0 \leq Y \leq K\} \neq \emptyset$ then there exists $\hat{\pi} \in S$ such that $\hat{\pi} \in \bar{f}(\hat{\pi})$.

Proof. First we prove f is upper hemicontinuous and then use this to prove \bar{f} is upper hemicontinuous; finally we apply the famous Kakutani fixed point theorem to show the existence of fixed point of \bar{f} .

For all $\hat{\pi}^i \in S$, $i = 1, 2, \dots$ and $\hat{\pi}^i \rightarrow \hat{\pi}$, for all $\pi^{*i} \in f(\hat{\pi}^i)$ and $\pi^{*i} \rightarrow \pi^*$, we show that $\pi^* \in f(\hat{\pi})$.

By our definition of f , for all $\pi^{*i} \in f(\hat{\pi}^i)$, there exists σ^{*i} , such that (σ^{*i}, π^{*i}) is an optimal solution of $(D\hat{\pi}^i)$. From strong duality theorem of linear programming, there exists (Y^{*i}, θ^{*i}) , an optimal solution of $(P\hat{\pi}^i)$, such that

$$\pi^{*i} b - \sigma^{*i} K = -\theta^{*i} . \quad (1)$$

Since (Y^{*i}, θ^{*i}) is feasible for $(P\hat{\pi}^i)$, $\theta^{*i}(d^0 - M\hat{\pi}^i) \leq AY^{*i} - b$, $i = 1, 2, \dots$ and by assumption, $d^0 - M\hat{\pi}^i \geq \bar{d} > 0$ $i = 1, 2, \dots$, $0 \leq Y \leq K$, we conclude that $\{\theta^{*i}, i = 1, 2, \dots\}$ is bounded. Combine this with (1), we know $\{\sigma^{*ik}, i = 1, 2, \dots\}$ is bounded. Since $K > 0$, $\sigma^{*i} \geq 0$, this implies $\{\sigma^{*i}, i = 1, 2, \dots\}$ is bounded. Therefore there exists a subsequence σ^{*ij} converges to σ^* . Without loss of generality, assume $\sigma^{*i} \rightarrow \sigma^*$, then from (1), we know that $\theta^{*i} \rightarrow \theta^*$, and

$$\pi^* b - \sigma^* K = -\theta^* . \quad (2)$$

Since (σ^{*i}, π^{*i}) is feasible for $(D\hat{\pi}^i)$, we have: $\pi^{*i} A \leq \sigma^{*i}$, $\pi^{*i}(-d^0 + M\hat{\pi}^i) \leq -1$, $\pi^{*i} \geq 0$, $\sigma^{*i} \geq 0$. Letting $i \rightarrow \infty$ we get $\pi^* A \leq \sigma^*$, $\pi^*(-d^0 + M\hat{\pi}) \leq -1$, $\pi^* \geq 0$, $\sigma^* \geq 0$, i.e., (σ^*, π^*) is feasible for $(D\hat{\pi})$. Similarly, we can assume $Y^{*i} \rightarrow Y^*$ and (Y^*, θ^*) is feasible

for $(P\hat{\pi})$. Because (2) holds, by weak duality theorem of linear programming, (Y^*, θ^*) is an optimal solution of $(P\hat{\pi})$, and (σ^*, π^*) is an optimal solution of $(D\hat{\pi})$, therefore, $\pi^* \in f(\hat{\pi})$, f is upper hemicontinuous.

Next, we prove \bar{f} is upper hemicontinuous.

For all $\hat{\pi}^i \in S$, $i = 1, 2, \dots$ and $\hat{\pi}^i \rightarrow \hat{\pi}$, for all $(\pi^{*i}/e\pi^{*i}) \in \bar{f}(\hat{\pi}_1)$ and $(\pi^{*i}/e\pi^{*i}) \rightarrow \pi^*$, by assumption $\{Y : AY > b, 0 \leq Y \leq K\} \neq \emptyset$, we know $\theta^{*i} > 0$, $i = 1, 2, \dots$, therefore by complement slackness, $\pi^{*i}(d^0 - M\hat{\pi}^i) = 1$, $i = 1, 2, \dots$. Since we assume $d^0 - M\hat{\pi}^i \geq \bar{d} > 0$, $i = 1, 2, \dots$, we know $\{\pi^{*i}, i = 1, 2, \dots\}$ is bounded. Without loss of generality, assume $\pi^{*i} \rightarrow \bar{\pi}$, then

$$\hat{\pi}^{*i}/e\pi^{*i} \rightarrow \bar{\pi}/e\bar{\pi} = \pi^*, \bar{\pi} = e\bar{\pi} \cdot \pi^*.$$

Since we have already shown that f is upper hemicontinuous, we know $\bar{\pi} = e\bar{\pi} \cdot \pi^* \in f(\hat{\pi})$, but $e\pi^* = 1$, therefore $\pi^* \in \bar{f}(\hat{\pi})$, therefore, \bar{f} is upper hemicontinuous.

Under the assumption of Theorem 1, the assumption of Lemma 1 still holds, so for all $\hat{\pi} \in S$, $\bar{f}(\hat{\pi})$ is a nonempty convex subset of S , and we have proved \bar{f} is upper hemicontinuous, therefore by Kakutani Fixed Point Theorem, \bar{f} has a fixed point. Q.E.D.

Next we weaken the assumption of Theorem 1, and prove the existence of a fixed point of \bar{f} .

Theorem 2. If $\{Y : AY \geq b, 0 \leq Y \leq K\} \neq \emptyset$, and for all $\hat{\pi} \in S$, $d^0 - M\hat{\pi} > 0$, $\|d^0 - M\hat{\pi}\| > \varepsilon > 0$, then there exists a $\hat{\pi} \in S$ such that $\hat{\pi} \in \bar{f}(\hat{\pi})$.

Proof. Since the assumption of Lemma 1 still holds, we know that for all $\hat{\pi} \in S$, $f(\hat{\pi}) \neq \emptyset$ and $\bar{f}(\hat{\pi})$ is a convex subset of S . The proof of upper hemicontinuity of f is similar to the proof in Theorem 1, so we leave it out. We now show that f is upper hemicontinuous implies \bar{f} is upper hemicontinuous.

For all $\hat{\pi}^i \in S$, $i = 1, 2, \dots$ and $\hat{\pi}^i \rightarrow \hat{\pi}$, for all $(\pi^{*i}/e\pi^{*i}) \in \bar{f}(\hat{\pi}^i)$ and $(\pi^{*i}/e\pi^{*i}) \rightarrow \pi^*$.

Case 1. If there exists a subsequence $\pi^{*ij} \rightarrow k\pi^*$ where $k \in (0, \infty)$, from the upper hemicontinuity of f , we know $k\pi^* \in f(\hat{\pi})$, but $\pi^* = k\pi^*/(k\pi^*) \in \bar{f}(\hat{\pi})$, therefore in this case \bar{f} is upper hemicontinuous.

Case 2. If for all $k \in (0, +\infty)$, there does not exist $\pi^{*ij} \rightarrow k\pi^*$, then there exists a subsequence π^{*ij} , $j = 1, 2, \dots$, such that $e\pi^{*ij} \rightarrow +\infty$. Without loss of generality, assume $e\pi^{*i} \rightarrow +\infty$. Since $\pi^{*i}b - \sigma^{*i}K = -\theta^{*i}$ still holds, $\{\theta^{*i}, i = 1, 2, \dots\}$ still bounded. Dividing the above equality by $e\pi^{*i}$, $(\pi^{*i}b/e\pi^{*i}) - (\sigma^{*i}K/e\pi^{*i}) = -(\theta^{*i}/e\pi^{*i})$. Letting $i \rightarrow \infty$, we have $\pi^{*i}b - \lim_{i \rightarrow \infty}(\sigma^{*i}K/e\pi^{*i}) = 0$. Letting $\xi^i = (\sigma^{*i}/e\pi^{*i})$, $\{\xi^i, i = 1, 2, \dots\}$ is bounded. Without loss of generality, assume $\xi^i \rightarrow \xi$, then $\xi \geq 0$ and $\lim_{i \rightarrow \infty}(\sigma^{*i}K/e\pi^{*i}) = \xi \cdot K$. Dividing $\pi^{*i}A \leq \sigma^{*i}$ and $\pi^{*i}(-d^0 + \hat{M}^i) \leq -1$ by $e\pi^{*i}$ and letting $i \rightarrow \infty$, we get $\pi^*A \leq \xi$, $\pi^*(-d^0 + \hat{M}) \leq 0$. Because $\pi^* \neq 0$ and $(-d^0 + \hat{M}) < 0$, there exists $k > 0$, such that $k\pi^*(-d^0 + \hat{M}) < -1$, $k\pi^*A \leq k\xi$, $k\pi^* \geq 0$, $k\xi \geq 0$, $k\pi^*b - k\xi \cdot K = 0$. This means $(k\xi, k\pi^*)$ is

feasible for $(D\hat{\pi})$. On the other hand, $\{Y : AY \geq b, 0 \leq Y \leq K\} \neq \emptyset$, so there exists Y^* , such that $(Y^*, 0)$ is feasible for $(P\hat{\pi})$. Because $k\pi^*b - k\xi^* \cdot K = 0$, by weak duality theorem, $(k\xi^*, k\pi^*)$ is an optimal solution of $(D\hat{\pi})$ and $(Y^*, 0)$ is an optimal solution of $(P\hat{\pi})$, therefore $k\pi^* \in f(\hat{\pi})$, $\pi^* = [k\pi^*/e(k\pi^*)] \in \bar{f}(\hat{\pi})$, i.e., \bar{f} is upper hemicontinuous. By Kakutani Fixed Point Theorem, \bar{f} has a fixed point. Q.E.D.

Economic Interpretation:

Let A be the technology matrix of an economy and $Y \geq 0$ the level of production, $Y = (Y_1, Y_2, \dots, Y_n)^T$. The net production available for consumption is AY . Let K be vector of capacities available so that $Y \leq K$. Consumption is a vector $b + \theta d$ where b is the fixed part, θ is a scalar, and d is the variable part that depends on relative prices. We assume, at fixed relative prices, the economy acts so as to maximize θ ,

$$\begin{aligned} \max \quad & \theta \\ \text{s.t.} \quad & AY - (b + \theta d) \geq 0 \\ & 0 \leq Y \leq K \end{aligned}$$

We assume the variable demand d depends on relative prices π . Thus if $\pi = \hat{\pi}$, let us suppose $d = -(1/e\hat{\pi}) M\hat{\pi}$, where M is positive definite but not necessarily symmetric matrix.

We play a little game. We guess values for $\hat{\pi}$, compute d , solve the LP and determine optimal θ^* , Y^* , σ^* , π^* . Next we form (normalized π^*) $\pi^*/e\pi^*$ and compare it with (normalized $\hat{\pi}$) $\hat{\pi}/e\hat{\pi}$. If equal, the game is over, i.e., we have found an equilibrium price. If not, we guess

another $\hat{\pi}$ and try again. Question is: is there a choice of $\hat{\pi}$ such that $(\pi^*/e\pi^*) = (\hat{\pi}/e\hat{\pi})$? Under the assumption of Theorem 1 or Theorem 2, such equilibrium prices exist.

Finally we would like to point out that the theorems and proofs can be generalized to a n-period planning model.

Acknowledgement:

The author would like to express her appreciation to Professor George B. Dantzig who suggested the problem and provided guidance throughout the research.

References:

1. G.B. Dantzig, Linear Programming and Extensions, Princeton University Press, Princeton, 1967.
2. C.B. Garcia, and W.I. Zangwill, Pathways to Solutions, Fixed Points, and Equilibria, Prentice Hall, Englewood Cliffs, 1981.
3. R.W. Cottle, Lecture notes on mathematical programming at Department of Operations Research, Stanford University.
4. B.C. Eaves, Lecture notes on mathematical programming at Department of Operations Research, Stanford University.

SOL 85-10: EXISTENCE OF EQUILIBRIUM PRICES FOR A SIMPLE PLANNING MODEL
by Hui Hu

Consider parametric LP:

$$\begin{array}{ll} \min & -\theta \\ \text{s.t.} & AY + \theta(-d^0 + M\hat{\pi}) \geq b \\ & 0 \leq Y \leq K, \quad \theta \geq 0 \end{array} \quad (I)$$

where M is a positive definite (not necessarily symmetric) matrix,
 $K > 0$, $\hat{\pi}$ is a parameter, $\hat{\pi} \in S = \{\pi \geq 0: e\pi = 1\}$.

For each fixed $\hat{\pi} \in S$, we can solve (I) and it's dual problem and get optimal θ^* , Y^* , σ^* , π^* . The question is, is there a $\hat{\pi} \in S$ such that after solving the corresponding LP and normalizing the dual price π^* , it turns out that $\pi^*/e\pi^* = \hat{\pi}$? In this paper, we are going to show that under certain conditions, such a $\hat{\pi}$ does exist.

The economic interpretation of the above model will be given at the end of the paper.

END

FILMED

9-85

DTIC